



**80th International Scientific
Conference of the
University of Latvia 2022**

CHEMISTRY SECTION

BOOK OF ABSTRACTS



**UNIVERSITY
OF LATVIA**

Reports	69
SYNTHESIS OF 2-AMINOQUINAZOLIN-4(3H)-ONE BASED PLASMEPSIN INHIBITORS	69
Emīls Edgars Basens	
SYNTHESIS OF BETA-ADRENERGIC RECEPTOR AGONISTS FOR THE TREATMENT OF OBESITY	70
Māris Bazulis	
COBALT-CATALYZED AMINO ACID C(sp ²)-H BOND FUNCTIONALIZATION USING ORGANIC ISOCYANIDES	71
Aleksandrs Čižikovs	
A MECHANISTICAL STUDY OF LEWIS BASE CATALYZED CYANOHYDRIN SYTHESIS	72
Artūrs Raimonds Feldmanis, Oto Filipsons	
SYNTHETIC APPROACH TOWARDS ENANTIOPURE CYCLIC SULFINAMIDES	73
Glebs Jersovs	
SYNTHESIS OF PURINE-THIAZOLOPYRIMIDINE CONJUGATES	74
Kārlis Ēriks Kriķis, Zigfrīds Kapilinskis	
C-H ACTIVATION OF LUPANE TYPE TRITERPENOIDs	75
Vladislavs Kroškins, Jevgeņija Lugiņina, Kristiāns Jankovičs	
AZIDE-TETRAZOLE EQUILIBRIUM IN PYRIDO[2,3-D]PYRIMIDINES	76
Kristaps Leškovskis	
SYNTHESIS OF BICYCLIC PROLINE ANALOGUES	77
Mērija Meiberga	
DEVELOPMENT OF SYNTHESIS PATHWAYS FOR THE LIMONOID OCTAHYDRO-1H-2,4-METHANOINDENE SCAFFOLD	78
Olavs Rāciņš	
THIAZOLINE CARBENE-Cu(I)-CARBAZOLIDE COMPLEXES AS LUMINESCENT TADF MATERIALS	79
Armands Rudušs, Žanis Sisojevs, Annija Jēce	
NEW SYNTHETIC APPROACH FOR THE SYNTHESIS OF SUBSTITUTED 7-ARYLPURINES	80
Armands Sebris	
COPPER(I) CATALYZED AZIDE-ALKYNE CYCLOADDITION IN IONIC LIQUIDS	81
Dīāna Sloboda	
SYNTHESIS OF 2-AMINOQUINAZOLINES AND INDAZOLES FROM 2-FORMYLPHENYLBORONIC ACIDS	82
Vitalii Solomin	


NEW SYNTHETIC APPROACH FOR THE SYNTHESIS OF SUBSTITUTED 7-ARYLPURINES

Armands Sebris

*Institute of Technology of Organic Chemistry, Faculty of Materials Science and Applied Chemistry, Riga Technical University, P. Valdena iela 3, Riga, Latvia
E-mail: armands.sebris_1@rtu.lv*

Alkylation and arylation on the purine ring usually proceeds almost selectively at N(9) position. While there are some simple pathways to introduce alkyl substituents at N(7) [1], the same is not true for introduction of aryl groups. The most commonly used Cu catalyzed Chan-Lam reaction [2] and arylation with iodanes [3] selectively give N(9) product. The few existing methods for purine N(7) arylation still provide a mixture of two isomers and are substrate dependent [4]. Hence, we decided to test various pathways towards 7-arylpurines starting from substituted pyrimidines.

We have achieved the best results with the following synthetic pathway starting from compound **1**: arylation was performed using diaryliodonane in the presence of Cu catalyst, giving 5-arylamino substituted pyrimidine **2**. Next, in S_NAr reaction under optimized conditions only one chlorine atom was substituted. Finally 7-arylpurine was obtained in a ring closing reaction with orthoester under acidic conditions. Modifications of starting material **1**, diaryliodonane reagent and ring closing reagent can be made to achieve differently substituted 7-arylpurines of type **4**.



Supervisors: Dr. chem. M. Turks, Dr. chem. I. Novosjolova.

Acknowledgment. This work was supported by the Latvian Council of Science grant No LZP-2020/1-0348.

References

- [1] Kotek, V.; Chudikova, N.; Tobrman, T.; Dvorak, D. *J. Org. Lett.*, **2010**, *12*, 5724.
- [2] Traskovskis, K.; Sebris, A.; Novosjolova, I.; Turks, M.; Guzauskas, M.; Volyniuk, D.; Bezikonnyi, O.; Grazulevicius, J. V.; Mishnev, A.; Grzibovskis, R.; Vembris, A. *J. Mater. Chem. C*, **2021**, *9*, 4532.
- [3] Niu H.-Y.; Xia, C.; Qu, G.-R.; Zhang, Q.; Jiang, Y.; Mao, R.-Z.; Li, D.-Y.; Guo, H.-M. *Org. Biomol. Chem.*, **2011**, *9*, 5039.
- [4] Keder, R.; Dvorakova, H.; Dvorak, D. *Eur. J. Org. Chem.* **2009**, *10*, 1522.